


# *Combined Serological Detection of Circulating Angiostrongylus vasorum Antigen and Parasite-specific Antibodies in Dogs from Hungary*

**Manuela Schnyder, Roland Schaper, Zoltán Lukács, Sándor Hornok & Róbert Farkas**

**Parasitology Research**  
Founded as Zeitschrift für  
Parasitenkunde

ISSN 0932-0113  
Volume 114  
Supplement 1

Parasitol Res (2015) 114:145-154  
DOI 10.1007/s00436-015-4520-1



**Your article is published under the Creative Commons Attribution license which allows users to read, copy, distribute and make derivative works, as long as the author of the original work is cited. You may self-archive this article on your own website, an institutional repository or funder's repository and make it publicly available immediately.**

# Combined Serological Detection of Circulating *Angiostrongylus vasorum* Antigen and Parasite-specific Antibodies in Dogs from Hungary

Manuela Schnyder<sup>1</sup>✉, Roland Schaper<sup>2</sup>, Zoltán Lukács<sup>3</sup>, Sándor Hornok<sup>4</sup>, Róbert Farkas<sup>4</sup>

<sup>1</sup> Institute of Parasitology, Vetsuisse Faculty, University of Zurich, 8057 Zurich, Switzerland

<sup>2</sup> Bayer Animal Health GmbH, 51368 Leverkusen, Germany

<sup>3</sup> Bayer Hungaria Kft., 1123 Budapest, Hungary

<sup>4</sup> Department of Parasitology and Zoology, Faculty of Veterinary Science Budapest, 1078 Budapest, Hungary

## Corresponding author:

Manuela Schnyder

✉ E-mail: manuela.schnyder@uzh.ch

## Abstract

The occurrence of the nematode *Angiostrongylus vasorum*, also known as the French heartworm, is increasingly being reported from various European countries. The adults of this parasite species live in the pulmonary arteries and right cardiac ventricle of wild canids and domestic dogs. Larval stages and eggs in the lungs induce inflammatory verminous pneumonia, causing severe respiratory disease in dogs. Furthermore, haematological and neurological signs and even death may occur. In Hungary, *A. vasorum* has been identified in red foxes, golden jackals and in two dogs and some slugs. In this first large-scale survey, 1247 sera from pet dogs were collected and tested by an ELISA for the detection of circulating antigen of *A. vasorum* and by a separate

ELISA to detect specific antibodies against the parasite. A total of 1.36% (n=17, 95% confidence intervals, CI: 0.80–2.17%) of the animals were positive in both ELISAs, while 1.76% (n=22, CI: 1.11–2.66%) of the tested dogs were antigen-positive only and 2.73% (n=34, CI: 1.90–3.79%) were positive for specific antibodies only. Regions with antigen- and antibody-positive animals overlapped and were distributed over nearly the whole sampled areas of the country. A considerable number of cases was observed in Budapest and also in the southern part of the country bordering Croatia, while in the most eastern part bordering Ukraine no positive samples were detected. These results confirm the endemic occurrence of *A. vasorum* in dogs originating from different parts of Hungary and the significant advantages of *A. vasorum* serology in epidemiological studies.

## Introduction

Canine angiostrongylosis is a potentially fatal disease caused by the metastrongylid nematode *Angiostrongylus vasorum*, which is increasingly being reported throughout Europe. The adult parasite lives in the pulmonary arteries and right cardiac ventricle of dogs, foxes and other wild carnivores, which are infected through the ingestion of obligatory intermediate hosts (snails or slugs) (Guilhon and Bressou 1960; Guilhon and Cens 1973) or paratenic hosts (Bolt et al. 1993) containing the infectious third stage larvae. Clinical signs in dogs most frequently include respiratory signs such as coughing and dyspnoea, but a broad range of further signs indicating coagulopathies or neurological dysfunctions (Chapman et al. 2004; Staebler et al. 2005; Wessmann et al. 2006; Koch and Willesen 2009), may be the signs most obvious to clinicians and animal owners. This variability and the fact that the disease is often in an advanced stage when noticed by the animal owners make the diagnosis of *A. vasorum* infections challenging but important: the sooner an appropriate anthelmintic treatment is initiated, the better for the clinical recovery of the dogs and limitation of damage, especially to lung tissues (Dennler et al. 2011; Schnyder et al. 2009).

A fundamental role in the early diagnosis of infections is played by awareness of the local occurrence of the parasite and, correspondingly, disease awareness among vets and animal owners. The endemic occurrence of *A. vasorum* is obviously linked to the presence of final hosts, among which red foxes represent an important wild reservoir, and the intermediate hosts, i.e. a broad range of snail and slug species (Eckert and Lämmler 1972; Ferdushy et al. 2009; Patel et al. 2014). Interestingly, although both final and intermediate hosts are widely present, the distribution of foxes and dogs affected by *A. vasorum* is scattered (Morgan 2014), and the reasons for this patchy distribution are not yet fully understood. Accordingly, it is particularly important to have good knowledge of the occurrence of

*A. vasorum* in a specific catchment area of a veterinary practice. Southern France (Guilhon and Cens 1969; Bourdeau 1993), Ireland (Dodd 1973), south-east England and Wales (Jacobs and Prole 1975; Simpson and Neal 1982) and Denmark (Bolt et al. 1992) have traditionally been endemic foci of *A. vasorum*; in the past two decades there has additionally been a sharp increase in reports from other countries in central Europe (Staebler et al. 2005; Barutzki and Schaper 2009), the Iberian peninsula (Segovia et al. 2004; Manas et al. 2005; Alho et al. 2014;) and also from Italy (Della Santa et al. 2002; Guardone et al. 2013), Greece (Papazahariadou et al. 2007) and Croatia (Rajkovic-Janje et al. 2002). Very recent case reports from eastern European countries describe the presence of the parasite in dogs from Poland (Demiaszkiewicz et al. 2014) and Slovakia (Miterpakova et al. 2014). In Hungary, which has common borders with Slovakia and Croatia, *A. vasorum* was first identified in 5 out of 100 red foxes shot in the western and southern counties of the country (Sréter et al. 2003), in two golden jackals (*Canis aureus*) (Takács et al. 2013) and also in two dogs and some slugs from close to the Croatian border diagnosed as being positive (Majoros et al. 2010).

The long-standing method for confirmation of clinical suspicion of dogs infected with *A. vasorum*, once the dogs are patent at approximately 40–57 days post infection (Guilhon and Cens 1973; Schnyder et al. 2010), is the detection of first stage larvae in faeces performed with the Baermann-Wetzel technique (Deplazes et al. 2013). However, by the time dogs start to be Baermann-positive, damage to the lung parenchyma is already present (Dennler et al. 2011; Guilhon and Cens 1969; Neff 1971). In addition, copromicroscopic methods have limitations concerning sensitivity and specificity due to variable larval excretion and the presence of other lungworm larvae (e.g. of *Crenosoma vulpis*) that may be excreted. Among the more recently applied techniques such as PCR (Jefferies et al. 2009; Al-Sabi et al. 2010) and serological methods (Cury et al. 1996; Verzberger-Epshtein et al. 2008) for

**Table 1:** Serological results of 1247 dog samples from Hungary tested for the presence of circulating antigens of *A. vasorum* and of specific antibodies against *A. vasorum*.

	Positive samples (n)	%	95% confidence intervals
Antibody-positive	51	4.09	3.06–5.34
Antibody-positive only	34	2.73	1.90–3.79
Antigen-positive	39	3.13	2.23–4.25
Antigen-positive only	22	1.76	1.11–2.66
Antibody- and antigen-positive	17	1.36	0.80–2.17

the detection of infected animals, serological methods have been shown to be highly suitable for both individual and population studies (Guardone et al. 2013; Schnyder et al. 2013a). Notably, mass screening with ELISAs is a valid and affordable method for investigation of large dog populations.

The aim of this study was to perform a seroepidemiological survey for the detection of circulating antigens of *A. vasorum* and of specific antibodies in sera of dogs originating from Hungary in order to increase knowledge of the presence of *A. vasorum* in dogs from this country.

## Materials and methods

Sera of 1247 dogs from Hungary were collected from dogs attending veterinary clinics for different reasons. Subsamples were sent to the Institute of Parasitology, Vetsuisse Faculty, University of Zurich, Switzerland and were analysed for the presence of circulating *A. vasorum* antigens using monoclonal and polyclonal antibodies in a sandwich ELISA with a sensitivity of 95.7% and a specificity of 94.0%, as previously described (Schnyder et al. 2011). Additionally, a sandwich ELISA (sensitivity 81.0%, specificity 98.8%) using *A. vasorum* adult somatic antigen purified by monoclonal antibodies (mAb Av 5/5) was used for specific antibody detection (Schucan et al. 2012). Test thresholds were regionally determined with 300 randomly selected samples based on the mean value of optical density ( $A_{405\text{ nm}}$ ) plus 3 (antigen detection) or 4 (antibody


detection) standard deviations. All test runs included a background control, a conjugate control, three positive control sera from three experimentally infected dogs and two negative control sera from uninfected dogs.

The collected data were analysed by a geographical information system (GIS) using the program RegioGraph 10 (GfK GeoMarketing, Bruchsal, Germany) to visualize the regional distribution of collected and analysed serum samples and *A. vasorum* antigen- and/or antibody-positive samples. The locations of positive samples were displayed on maps with administrative and postcode boundaries based on the three-digit postcodes of Hungary as points of reference.

Excel 2007 for Windows (Microsoft Corporation, Redmond, USA) was used to calculate the prevalence values and the 95% confidence interval (CI) of prevalence values.

## Results

The seropositivity of all tested samples is summarised in Table 1. A total of 1.36% (n=17, confidence intervals, CI: 0.80–2.17%) of the animals were positive in both ELISAs, while 1.76% of the tested dogs were only antigen-positive and 2.73% were positive for specific antibodies only. The locations of positive sera are shown in Fig. 1a–c. Regions with antigen- and antibody-positive animals overlapped and were distributed over nearly the whole sampled areas of the country. The area of Budapest is



**Fig. 1:** Occurrence of *Angiostrongylus vasorum* in 1247 dogs from Hungary detected by ELISA.

Dark grey areas represent the origin of the tested dog sera.

**Fig. 1a:** Dogs positive for detection of specific antibodies against *A. vasorum*.

enlarged, showing a considerable number of cases. An accumulation of cases can also be observed in the southern part of the country bordering Croatia, while in the most eastern part bordering Ukraine no positive samples were detected.

## Discussion

With 1.36% of the examined dogs being positive in both ELISAs, the prevalence in Hungary is significantly higher than that found for Germany (Schnyder et al. 2013a), Italy (Guardone et al. 2013) or Poland (Schnyder et al. 2013b), which were all between 0.3 and 0.5% and all obtained using the same validated procedures. More than 4% of the dogs were positive for specific antibodies

against *A. vasorum*, indicating parasite exposure: these dogs may have been sampled during the first five weeks after an *A. vasorum* infection, when antigen detection is still negative, or the dogs were parasite-free but still antibody-positive after anthelmintic treatment or natural clearance of the infection (Schnyder et al. 2013a). The procedures used, i.e. the combination of both ELISAs, again confirmed their utility for mass screening of dog populations. In contrast to Poland, where testing of more than 3000 sera revealed for the first time the presence of *A. vasorum* in the country (Schnyder et al. 2013b) and positive foxes were identified subsequently (Demiaszkiewicz et al. 2014), the occurrence of *A. vasorum* in Hungary had already been reported. The parasite had first been described in foxes: 5 of 100 red foxes (*Vulpes*




Fig. 1b: Dogs positive for detection of circulating antigen of *A. vasorum*.

*vulpes*) from 16 Hungarian counties dissected in 2002 were positive at necropsy, with four foxes having low worm burdens (1–7 adult specimens) and a single fox having a moderate worm burden (8–14 specimens). These foxes originated from the southern and western parts of the country (Sréter et al. 2003), while none of the surrounding countries had reported cases of *A. vasorum* at that stage. Two asymptomatic dogs from south-western Hungary were diagnosed positive for *A. vasorum* based on positive faecal samples (Majoros et al. 2010). These cases were followed more recently by further reports of *A. vasorum* in wild carnivores, for instance in 2 out of 10 golden jackals (*Canis aureus*) (Takács et al. 2013). In a current analysis of the environmental factors having an impact on the distribution of lung worms in foxes based on the dissection of 937 animals, an average prevalence

of 17.9% was detected, with high variations from 0–59.4% between counties (Tolnai et al. 2015). Interestingly, *A. vasorum* was nearly absent in the eastern and central-southern part of the country, while *C. vulpis* was also detected in the most eastern part. This latter lung worm species, which is located in the bronchi and bronchioles of dogs and foxes, is also a cause of respiratory diseases and uses snails and slugs as intermediate hosts (Stockdale and Hulland 1970; Unterer et al. 2002). Its prevalence in foxes in 2013/14 (24.6%) (Tolnai et al. 2015) was identical to that detected in 2002 (24%) (Sréter et al. 2003). Therefore, if the presence of *C. vulpis* is long-standing and stable in Hungary, data from infected foxes and dogs show that *A. vasorum* is a more recent phenomenon. The occurrence of canine angiostrongylosis is actually less frequent, and in foxes its dissemination is not




Fig. 1c: Combined testing for detection of *A. vasorum* antigen and specific antibodies.

as broad as for *C. vulpis*, although they may share the same intermediate hosts and mollusc species are broadly distributed in Hungary (Sólymos and Fehér 2005).

During the last few decades, reports of animals affected by *A. vasorum* animals have become more frequent and various reasons for the apparent spread of lung worms have been discussed. Regional climate changes in vector epidemiology and movements in animal populations were taken into account (Morgan et al. 2009; Tolnai et al. 2015). Dog data collected in this study support the gradual advancement of *A. vasorum* in eastern Europe: in Hungary the parasite has been detected in parts of the country bordering Croatia in the south and Slovakia in the north, where the parasite has been already reported (Miterpakova et al. 2014; Rajkovic-Janje et al. 2002), while little is known from the

other neighbouring countries (Austria, Romania, Slovenia, Serbia and Ukraine).

The evidence of a higher prevalence of all important lungworms (*A. vasorum*, *C. vulpis* and *Eucoleus aerophilus*) in foxes (Sréter et al. 2003; Tolnai et al. 2015) compared to dogs (Koch and Willeßen 2009) indicates that wildlife, in the absence of obvious geographical barriers, plays an important role in expansion and establishment of these parasites. Hungary, like many other countries, has implemented successful rabies control programmes (Solymosi et al. 2002), which implies a massive increase in the fox population. Therefore, it is of general acceptance that foxes contribute essentially to the local distribution and establishment of *A. vasorum*, while infected dogs that travel long distances may contribute to a broader spread beyond connected endemic areas.

The major clinical impact of *A. vasorum* infections in dogs means that it is of essential importance to increase awareness of the occurrence of *A. vasorum* among clinicians and dog owners, particularly in previously unreported areas bordering endemic regions, with the aim of preventing fatal cases of canine angiostrongylosis. Newly developed, rapid and easy-to-use commercial devices for use in veterinary practices (Schnyder et al. 2014) and adequate metaphylactic and prophylactic anthelmintic treatments (Conboy 2001; Willesen et al. 2007; Schnyder et al. 2009) allow early diagnosis and consequently a good prognosis for dogs affected by *A. vasorum*.

## References

- Al-Sabi MN, Deplazes P, Webster P, Willesen JL, Davidson RK, Kapel CM (2010) PCR detection of *Angiostrongylus vasorum* in faecal samples of dogs and foxes. *Parasitol Res* 107:135–140
- Alho AM, Schnyder M, Meireles J, Belo S, Deplazes P, Madeira de Carvalho L (2014) Preliminary results on the seroprevalence of *Angiostrongylus vasorum* and co-infection with *Dirofilaria immitis* in shelter dogs from Portugal. *Paras & Vectors* 7 Suppl 1:O26
- Barutzki D, Schaper R, (2009) Natural infections of *Angiostrongylus vasorum* and *Crenosoma vulpis* in dogs in Germany (2007–2009). *Parasitol Res* 105 Suppl 1: S39–48
- Bolt G, Monrad J, Henriksen P, Dietz HH, Koch J, Bindseil E, Jensen AL (1992) The fox (*Vulpes vulpes*) as a reservoir for canine angiostrongylosis in Denmark. Field survey and experimental infections. *Acta Vet Scand* 33: 357–362
- Bolt G, Monrad J, Henriksen P, Dietz HH, Koch J, Bindseil E, Jensen AL (1993) The common frog (*Rana temporaria*) as a potential paratenic and intermediate host for *Angiostrongylus vasorum*. *Parasitol Res* 79, 428–430
- Bourdeau P (1993) L'angiostrongylose canine. *Receuil Méd Vét* 169: 401–407
- Chapman PS, Boag AK, Guitian J, Boswood A (2004) *Angiostrongylus vasorum* infection in 23 dogs (1999–2002). *J Small Anim Practice* 45: 435–440
- Conboy G (2004) Natural infections of *Crenosoma vulpis* and *Angiostrongylus vasorum* in dogs in Atlantic Canada and their treatment with milbemycin oxime. *Vet Rec* 155, 16–18
- Cury MC, Lima WS, Vitor RWA (1996) Enzyme-Linked Immunosorbent Assay (ELISA) for the diagnosis of *Angiostrongylus vasorum* (Baillet, 1866) infection in dogs. *Revue Méd Vét* 147: 525–530
- Della Santa D, Citi D, Marchetti V, Nardoni S (2002) Infestazione da *Angiostrongylus vasorum* nel cane: review della letteratura e presentazione di un caso clinico. *Veterinaria* 16: 9–14
- Demiaszkiewicz AW, Pyziel AM, Kuligowska I, Lachowicz J (2014) The first report of *Angiostrongylus vasorum* (Nematoda; Metastrongyloidea) in Poland, in red foxes (*Vulpes vulpes*). *Acta Parasitol* 59: 758–762
- Dennler M, Makara M, Kranjc A, Schnyder M, Ossent P, Deplazes P, Ohlerth S, Glaus TM (2011) Thoracic computed tomography findings in dogs experimentally infected with *Angiostrongylus vasorum*. *Vet Radiol Ultrasound* 52: 289–294
- Deplazes P, Eckert J, Samson-Himmelstjerna, Gv, Zahner H (2013) *Lehrbuch der Parasitologie für die Tiermedizin* (Stuttgart, Enke Verlag).
- Dodd K (1973) *Angiostrongylus vasorum* (Baillet, 1866) infestation in a greyhound kennels. *Vet Rec* 92: 195–197

### Ethical standards

All institutional and national guidelines for the care and use of laboratory animals were followed.

**Funding:** The study was partially funded by Bayer Animal Health.

### Conflict of interest

M. Schnyder, S. Hornok and R. Farkas declare no conflict of interest, R. Schaper and Z. Lukács are employed by Bayer Animal Health.

### Acknowledgements

We are grateful to Kathrina Stebler and Katja Huggel from the Institute of Parasitology in Zurich and Mónica Gyurkovszky from the Department of Parasitology and Zoology at the Faculty of Veterinary Science in Budapest for their extremely valuable support.

- Eckert J, Lämmler G (1972) Angiostrongylose bei Mensch und Tier. *Parasitol Res* 39: 303–322
- Ferdusly T, Kapel CM, Webster P, Al-Sabi MN, Gronvold J (2009) The occurrence of *Angiostrongylus vasorum* in terrestrial slugs from forests and parks in the Copenhagen area, Denmark. *J Helminthol* 83: 379–383
- Guardone L, Schnyder M, Macchioni F, Deplazes P, Magi M (2013) Serological detection of circulating *Angiostrongylus vasorum* antigen and specific antibodies in dogs from central and northern Italy. *Vet Parasitol* 192: 192–198.
- Guilhon J, Bressou C (1960) Rôle des Limacidés dans le cycle évolutif d'*Angiostrongylus vasorum* (Baillet, 1866). *C R Acad Sc* 251: 2252–2253
- Guilhon J, Cens B (1969) [Migrations and evolution of *Angiostrongylus vasorum* (Baillet, 1866) in dogs (in French)]. *Comptes Rendus Hebdomadaires des Seances de l'Academie des Sciences Naturelles* 269: 2377–2380.
- Guilhon J, Cens B (1973) *Angiostrongylus vasorum* (Baillet, 1866): Étude biologique et morphologique. *Ann Parasitol Hum Comp* 48: 567–596
- Jacobs DE, Prole JH (1975) *Angiostrongylus vasorum* and other nematodes in British greyhounds. *Vet Rec* 96: 180
- Jefferies R, Morgan ER, Shaw SE (2009) A SYBR green real-time PCR assay for the detection of the nematode *Angiostrongylus vasorum* in definitive and intermediate hosts. *Vet Parasitol* 166: 112–118
- Koch J, Willeßen JL (2009) Canine pulmonary angiostrongylosis: An update. *Vet J* 179: 348–359
- Majoros G, Fukar O, Farkas R, (2010) Autochthonous infection of dogs and slugs with *Angiostrongylus vasorum* in Hungary. *Vet Parasitol* 174: 351–354.
- Manas S, Ferrer D, Castella J, Maria Lopez-Martin J (2005) Cardiopulmonary helminth parasites of red foxes (*Vulpes vulpes*) in Catalonia, northeastern Spain. *Vet J* 169: 118–120
- Miterpakova M, Hurnikova Z, Zalewski AP (2014) The first clinically manifested case of angiostrongylosis in a dog in Slovakia. *Acta Parasitol* 59: 661–665
- Morgan ER (2014) Canine pulmonary angiostrongylosis: can a worm change its spots? *Vet Rec* 175: 116–117
- Morgan ER, Jefferies R, Krajewski M, Ward P, Shaw SE (2009) Canine pulmonary angiostrongylosis: the influence of climate on parasite distribution. *Parasitol Int* 58: 406–410
- Neff H, (1971) Experimentelle Infektionen von Hunden mit *Angiostrongylus vasorum* (Nematoda). Dissertation, Universität Zürich
- Papazahariadou A, Founta A, Papadopoulos E, Chliounakis S, Antoniadou-Sotiriadou K, Theodorides Y (2007) Gastrointestinal parasites of shepherd and hunting dogs in the Serres Prefecture, Northern Greece. *Vet Parasitol* 148: 170–173.
- Patel Z, Gill AC, Fox MT, Hermosilla C, Backeljau T, Breugelmans K, Keevash E, McEwan C, Aghazadeh M, Elson-Riggins JG (2014) Molecular identification of novel intermediate host species of *Angiostrongylus vasorum* in Greater London. *Parasitol Res* 113: 4363–4369
- Rajkovic-Janje R, Marinculic A, Bosnic S, Benic M, Vinkovic B, Mihaljevic Z (2002) Prevalence and seasonal distribution of helminth parasites in red foxes (*Vulpes vulpes*) from the Zagreb County (Croatia). *Z Jagdwiss* 48: 151–160
- Schnyder M, Fahrion A, Ossent P, Kohler L, Webster P, Heine J, Deplazes P (2009) Larvicidal effect of imidacloprid/moxidectin spot-on solution in dogs experimentally inoculated with *Angiostrongylus vasorum*. *Vet Parasitol* 166: 326–332
- Schnyder M, Fahrion A, Riond B, Ossent P, Webster P, Kranjc A, Glaus T, Deplazes P (2010) Clinical, laboratory and pathological findings in dogs experimentally infected with *Angiostrongylus vasorum*. *Parasitol Res* 107: 1471–1480
- Schnyder M, Tanner I, Webster P, Barutzki D, Deplazes P (2011) An ELISA for sensitive and specific detection of circulating antigen of *Angiostrongylus vasorum* in serum samples of naturally and experimentally infected dogs. *Vet Parasitol* 179: 152–158.
- Schnyder M, Schaper R, Bilbrough G, Morgan ER, Deplazes P, (2013a) Seroepidemiological survey for canine angiostrongylosis in dogs from Germany and the UK using combined detection of *Angiostrongylus vasorum* antigen and specific antibodies. *Parasitology* 140: 1442–1450
- Schnyder M, Schaper R, Pantchev N, Kowalska D, Szwedko A, Deplazes P, (2013b) Serological detection of circulating *Angiostrongylus vasorum* antigen- and parasite-specific antibodies in dogs from Poland. *Parasitol Res* 112 Suppl 1: 109–117
- Schnyder M, Stebler K, Naucke TJ, Lorentz S, Deplazes P (2014) Evaluation of a rapid device for serological in-clinic diagnosis of canine angiostrongylosis. *Paras & Vectors* 7: 72
- Schucan A, Schnyder M, Tanner I, Barutzki D, Traversa D, Deplazes P (2012) Detection of specific antibodies in dogs infected with *Angiostrongylus vasorum*. *Vet Parasitol* 185: 216–224.

- Segovia JM, Torres J, Miquel J (2004) Helminth parasites of the red fox (*Vulpes vulpes* L., 1758) in the Iberian Peninsula: an ecological study. *Acta Parasitologica* 49: 67–79
- Simpson VR, Neal C (1982) *Angiostrongylus vasorum* infection in dogs and slugs. *Vet Rec* 111: 303–304
- Sólymos P, Fehér Z (2005) Conservation prioritization based on the distribution of land snails in Hungary. *Conserv Biol* 19: 1084–1094
- Solymosi N, Reiczigel J, Edelyi K, Banhidly J, Foldi Z, Bodis K (2002) Spatial analysis of rabies cases in foxes in Hungary between 1990 and 2001: a preliminary report. *Stud Health Technol Inform* 90: 770–773
- Sréter T, Széll Z, Marucci G, Pozio E, Varga I (2003) Extraintestinal nematode infections of red foxes (*Vulpes vulpes*) in Hungary. *Vet Parasitol* 115: 329–334
- Staebler S, Ochs H, Steffen F, Naegeli F, Borel N, Sieber-Ruckstuhl N, Deplazes P (2005) Autochthonous infections with *Angiostrongylus vasorum* in dogs in Switzerland and Germany (in German). *Schweiz Arch Tierheilkde* 147:121–127
- Stockdale PH, Hulland TJ, (1970) The pathogenesis route of migration, and development of *Crenosoma vulpis* in the dog. *Pathol Vet* 7: 28–42
- Takács A, Szabo L, Juhász L, Takács AA, Lanszki J, Takács PT, Heltai M (2013) Data on the parasitological status of golden jackal (*Canis aureus* L., 1758) in Hungary. *Acta Vet Hung*: 1–9.
- Tolnai Z, Széll Z, Sréter T (2015) Environmental determinants of the spatial distribution of *Angiostrongylus vasorum*, *Crenosoma vulpis* and *Eucoleus aerophilus* in Hungary. *Vet Parasitol* 207: 355–358
- Unterer S, Deplazes P, Arnold P, Flückiger M, Reusch CE, Glaus TM (2002) Spontaneous *Crenosoma vulpis* infection in 10 dogs: laboratory, radiographic and endoscopic findings. *Schweiz Arch Tierheilkd* 144: 174–179
- Verzberger-Epshtein I, Markham RJ, Sheppard JA, Stryhn H, Whitney H, Conboy GA (2008) Serologic detection of *Angiostrongylus vasorum* infection in dogs. *Vet Parasitol* 151: 53–60
- Wessmann A, Lu D, Lamb CR, Smyth B, Mantis P, Chandler K, Boag A, Cherubini GB, Cappello R (2006) Brain and spinal cord haemorrhages associated with *Angiostrongylus vasorum* infection in four dogs. *Vet Rec* 158: 858–863
- Willesen JL, Kristensen AT, Jensen AL, Heine J, Koch J (2007) Efficacy and safety of imidacloprid/moxidectin spot-on solution and fenbendazole in the treatment of dogs naturally infected with *Angiostrongylus vasorum* (Baillet, 1866). *Vet Parasitol* 147:258–264.

